

Health & Ecological Risk Assessment

Conceptual Framework for Assessing Ecosystem Health

Mark A Harwell,^{*†} John H Gentile,[†] Larry D McKinney,[‡] John W Tunnell Jr,^{**‡} William C Dennison,[§] R Heath Kelsey,[§] Kiersten M Stanzel,^{||} Gregory W Stunz,[‡] Kim Withers,[‡] and Jace Tunnell[#]

[†]Harwell Gentile & Associates, LC, Port Orange, Florida, USA

[‡]Harte Research Institute for Gulf of Mexico Studies, Texas A&M University–Corpus Christi, Corpus Christi, Texas, USA

[§]University of Maryland Center for Environmental Science, Cambridge, Maryland, USA

^{||}Coastal Bend Bays & Estuaries Program, Corpus Christi, Texas, USA

[#]Mission-Aransas National Estuarine Research Reserve, Port Aransas, Texas, USA

ABSTRACT

Over the past century, the environment of the Gulf of Mexico has been significantly altered and impaired by extensive human activities. A national commitment to restore the Gulf was finally initiated in response to the unprecedented *Deepwater Horizon* oil spill in 2010. Consequently, there is a critical need for an assessment framework and associated set of indicators that can characterize the health and sustainability of an ecosystem having the scale and complexity of the Gulf. The assessment framework presented here was developed as an integration of previous ecological risk- and environmental management-based frameworks for assessing ecosystem health. It was designed to identify the natural and anthropogenic drivers, pressures, and stressors impinging on ecosystems and ecosystem services, and the ecological conditions that result, manifested as effects on valued ecosystem components. Four types of societal and ecological responses are identified: reduction of pressures and stressors, remediation of existing stressors, active ecosystem restoration, and natural ecological recovery. From this conceptual framework are derived the specific indicators to characterize ecological condition and progress toward achieving defined ecological health and sustainability goals. Additionally, the framework incorporates a hierarchical structure to communicate results to a diversity of audiences, from research scientists to environmental managers and decision makers, with the level of detail or aggregation appropriate for each targeted audience. Two proof-of-concept studies were conducted to test this integrated assessment and decision framework, a prototype Texas Coastal Ecosystems Report Card, and a pilot study on enhancing rookery islands in the Mission-Aransas Reserve, Texas, USA. This Drivers–Pressures–Stressors–Condition–Responses (DPSCR₄) conceptual framework is a comprehensive conceptual model of the coupled human–ecological system. Much like its predecessor, the ecological risk assessment framework, the DPSCR₄ conceptual framework can be tailored to different scales of complexity, different ecosystem types with different stress regimes, and different environmental settings. *Integr Environ Assess Manag* 2019;15:544–564. © 2019 The Authors. *Integrated Environmental Assessment and Management* published by Wiley Periodicals, Inc. on behalf of Society of Environmental Toxicology & Chemistry (SETAC)

Keywords: Ecological indicators Ecosystem health Ecosystem restoration Human–ecological system Integrated assessments

INTRODUCTION

We present here a conceptual framework with broad applicability for characterizing human–ecosystem interactions and for assessing ecosystem health. This initiative

began with the objective of constructing a framework for assessment of the health of the Gulf of Mexico; our discussion, therefore, begins with that rationale. However, as the conceptual framework evolved, it became clear that its applicability is not limited to the Gulf of Mexico and its utility is not limited to a report card on progress toward desired conditions. Rather, we present this conceptual framework both as an organizing guide for assessing the health of an ecosystem, the forces that affect it, and potential management avenues to achieve defined goals, and as a generic conceptual model of the coupled human–ecological system, adaptable to virtually any ecosystem and any environmental problem set.

**Deceased

This article includes online-only Supplemental Data.

* Address correspondence to mharwell@ecologicalrisk.com

Published 25 March 2019 on wileyonlinelibrary.com/journal/ieam.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial, and no modifications or adaptations are made.

The Gulf of Mexico is among the most ecologically diverse and valuable ecosystems in the world, comprising more than 1.5×10^6 km² in area and consisting of offshore waters and coastal habitats of 11 US and Mexican states plus Cuba. The Gulf's wetlands, beaches, coastal woodlands, and islands are major nurseries for breeding birds and provide foraging and stopover habitat for billions of birds that converge from some of the most important migratory flyways in the western hemisphere (Horton et al. 2019). Coastal marshes and nearshore habitats provide essential nursery habitat for ecologically, commercially, and recreationally important species of fish and invertebrates. Offshore habitats and species are biologically diverse and include deepwater corals, sponges, fish stocks, marine mammals, sea turtles, and other unique species and communities. These habitats are integral to the economic and cultural fabric of the Gulf, providing a range of ecosystem services, including fisheries, food and energy production, infrastructure protection, and recreational and wildlife-related activities. Testament to its impressive diversity is a recent biotic survey that found more than 15 400 species living in the Gulf of Mexico, including more than 2500 species of crustaceans, a similar number of mollusks, and 1975 species of vertebrates, comprising more than 1500 species of fish, 400 species of birds, and 30 species of mammals (Felder et al. 2009).

The Gulf's watershed covers more than half the continental USA (USEPA 2011), 40% from the Mississippi River Basin alone. This watershed is a source of a wide range of anthropogenic stressors: Nutrients (N and P) and other pollutants (e.g., hydrocarbons, pesticides, industrial wastes) contribute to degraded water quality in the Gulf, including more than 17 000 km² of annually occurring hypoxic conditions (USEPA 2011), peaking at 22 720 km² in 2017 (NOAA 2017). Oil and gas industry canals, pipelines, and other infrastructure crisscross the landscape, contributing to the loss of wetland habitat. Geologic land subsidence substantially exacerbates sea-level rise (Morton et al. 2005); for example, approximately 5000 km² of wetlands in Louisiana were lost in the last 7 decades (Couvillion et al. 2011). As a result of these and other natural and anthropogenic stressors, Gulf coastal ecosystems have become increasingly degraded for both human use and aquatic life. Scientific consensus has emerged for several continuing major threats to the health of the Gulf, including the following (Mabus 2010; USEPA 2011):

- loss of wetland habitats, coastal marshes, barrier islands, and shorelines;
- erosion of barrier islands and shorelines, undermining storm protection and reducing habitat for endangered or threatened species such as sea turtles and shorebirds;
- degradation of coastal estuaries, which provide essential nursery habitat for most of the Gulf fishery resources;
- overharvesting of commercially and recreationally important fisheries, exacerbated by the human health threats of methylmercury and other contaminants in finfish, harmful algal blooms (HABs), and human pathogens in shellfish;

- a large dead zone created by hypoxia offshore of the Mississippi River Delta; and
- global climate change with potentially increased frequency and intensity of storms, accelerated sea-level rise, and attendant economic risks and loss of coastal habitats and natural resources.

Superimposed on these threats was the April 2010 explosion on the *Deepwater Horizon* drilling platform operating in the Mississippi Canyon of the Gulf, resulting in the largest marine oil spill in US history, with an estimated 5×10^6 barrels released over 87 d (Mabus 2010; NAS 2012). The unprecedented combination of extreme depth of discharge (~1500 m) and massive use of dispersants ($\sim 3 \times 10^6$ L; Kujawinski et al. 2011) caused high uncertainty in predicting the transport and fate of oil and dispersant compounds and in understanding the severity and magnitude of ecological effects (Joye 2015).

In response to the oil spill, the Gulf Coast Ecosystem Restoration Task Force was established (Executive Order 13554, 5 October 2010) to develop a science-based Gulf of Mexico Regional Ecosystem Restoration Strategy to restore and conserve habitat, restore water quality, replenish and protect living coastal and marine resources, and enhance community resilience (USEPA 2011; <https://www.restorethegulf.gov>). This strategy requires an integrated, risk-based ecosystem assessment framework for informing decision making to achieve specific restoration goals. This framework in turn requires the identification of indicators and measures of success to evaluate the efficacy of the restoration program in meeting its goals. Indicators, along with measures of performance, must be quantifiable and understandable to the public, reflect the desired Gulf condition, and be sensitive to ecosystem changes (USEPA 2011; GCERC 2016).

We initiated a project to develop such an integrated assessment framework with associated set of indicators and metrics that could be used to characterize the health of the Gulf of Mexico ecosystems, including their linkages to human communities, termed the “Gulf EcoHealth Metrics Initiative.” Our vision was to develop a comprehensive conceptual framework for the coupled human–ecological system from which an assessment of the environmental condition of the Gulf could be derived that is scientifically based, widely accessible, and readily understandable by policy makers, stakeholders, scientists, and the American public. A hierarchical communications structure, unified by a common conceptual framework, provides the basis for informing multiple audiences at the appropriate level of detail and aggregation, allowing one to dig deeper into the reasons for the various assigned metrics of ecosystem health.

The ultimate aim of the Gulf EcoHealth initiative is to provide the scientific information and understanding necessary to evaluate the health of the Gulf, clearly demonstrate how well it is or is not progressing toward desired long-term goals, and inform the decision-making process on the policies and resources needed to achieve sustainability of a healthy Gulf of Mexico. As the initiative proceeded, however, it became

evident that although the assessment framework presented below (DPSCR₄ Framework: Need for a New Synthesis Framework) provides the conceptual foundation for achieving this goal, fully populating the elements of the framework may be more aspirational than feasible because of the insufficient availability of environmental data, both at present and in the plausible future. Nevertheless, this assessment framework can indicate what types of data would be most useful to have and can provide the rationale and priorities for designing an appropriate monitoring program. Its greater utility, however, is as a comprehensive conceptual framework for distinguishing those aspects of ecosystems and human–ecosystem interactions that matter from those that do not, thereby providing a roadmap for improved understanding of human interactions with ecosystems and a guide for environmental management decision making. Further, much like its predecessor, the ecological risk assessment (ERA) framework, this conceptual framework can be tailored to different scales of complexity, different ecosystem types with different stress regimes, and different environmental settings.

History of ecological health assessment frameworks

Environmental assessment indicators and report cards have become widespread as tools to characterize the status and trends of ecosystem health and to inform the allocation of resources for sustainability of healthy marine and coastal environments. The San Francisco Bay Index (The Bay Institute 2003), Gulf of Maine ecosystem indicators partnership (Mills 2006), Southeast Queensland healthy waterways report cards (Pantus and Dennison 2005), Chesapeake Bay Report Card (Williams et al. 2009, 2010; IAN 2013), Australia's Great Barrier Reef Report Card (Australian and Queensland Governments 2010), Florida Keys Ecosystem Report Card (NOAA 2011), scorecards for Marine Protected Areas (CEC 2011), US National Coastal Condition Report (USEPA 2012), Ocean Health Index (Halpern et al. 2012), the semiannual System-wide Ecological Indicators for Everglades Restoration reports (e.g., Brandt et al. 2016; www.evergladesrestoration.gov), and the America's Watershed Initiative (2015; americaswatershed.org) Mississippi River Watershed Report Card are examples of indicators and assessments being used to inform the public and decision makers about the health and sustainability of coastal ecosystems.

We reviewed the extensive literature on indicators of ecological condition and the literature on the conceptual frameworks for these and many other environmental assessments. Two approaches dominate, one derived from the perspective of stress ecology (e.g., Odum 1969, 1985; Holling 1973; Barrett et al. 1976; Rapport et al. 1985) and its derivatives, ERA and ecological indicators (e.g., Kelly and Harwell 1989, 1990; Gentile and Slimak 1990; USEPA 1992, 1998; Environment Canada 1994; Gentile et al. 1993; Harwell, Myers et al. 1999; Dale and Beyeler 2001; USEPA SAB 2002; Doren et al. 2009). In this approach, ecological condition or health is a result of causal stress–effect relationships, as manifested in specific indicators of selected

components (both structural and functional) of ecosystems. Here stressors are defined as physical, chemical, or biological agents that can cause effects on ecological systems. Effects are manifested as changes in specific ecological attributes that are ecologically and/or societally important, often termed “assessment endpoints” (USEPA 1998) or “valued ecosystem components” (VECs) (CCME 1996; Harwell et al. 2011). This approach seeks to elucidate the causal mechanisms of ecological effects from human activities and natural processes; consequently, it is closely related to hypothesis-driven scientific studies on how ecosystems and their components respond to environmental stressors, whether natural or anthropogenic. However, a limitation of this approach, particularly at larger scales, is that there may be too many environmental stressors to be managed, exacerbated by too many interactions among stressors and too many pathways leading to effects.

The second approach is more environmental-management focused, based on the Pressure–State–Response (PSR) framework (OECD 1991, 1993) and its derivative, the Drivers–Pressures–State–Impacts–Response (DPSIR) framework (EEA 1999; Weber 2010). In the latter, drivers are the fundamental forces causing pressures that affect the state of the environment; impacts are how the state changes because of the pressures; responses are societal feedbacks through adaptation or curative action. The pressures in DPSIR originally excluded natural processes except for climate change, but more recent applications have relaxed that exclusion (e.g., Weber 2010). The DPSIR framework has been adopted by the United Nations, European Union, and some US agencies because it is more attuned to the needs of decision makers, stakeholders, and the public when addressing environmental issues on large scales. However, a significant deficiency of the PSR and DPSIR approaches is that pressures are typically defined at such a broad level (e.g., population growth, agricultural production) that their relationships to the state of the environment are by necessity correlative instead of causal. Hence, it may provide insufficient specificity of the relationships between human activities and ecological effects to identify what needs to be managed and what management actions would be required in order to achieve a healthy environment. The other serious deficiency of DPSIR is its distinction between impacts and state, implying that there exists some baseline or natural state for an ecosystem and that impacts constitute some deviation from that state; characterized this way, human interactions are conceived of only as adverse and no accommodation is provided for the positive contributions of ecosystem services (Kelble et al. 2013). In reality, ecosystems are dynamic, changing over time and space, and ecosystems became fundamentally altered by human presence long ago, so there is no default baseline against which impacts are measured.

Irrespective of the assessment framework used, the indicators to measure environmental condition have also been a topic of considerable research and discussion over the past few decades. The literature on the utility and purposes for ecological indicators and criteria for selecting them was

developed early on (e.g., Kelly and Harwell 1989, 1990; Gentile and Slimak 1990; Hunsaker and Carpenter 1990; MacKenzie et al. 1990; Cairns et al. 1993). Some publications proposed specific indicators or indices that have been widely adopted, such as Karr’s (1981) fish community-based index of biotic integrity to characterize the condition of freshwater streams, and the Landres et al. (1988) use of vertebrate indicator species to characterize wildlife habitat quality. Other ecological indicators have been suggested from the molecular (e.g., Goksøyr and Förlin 1992) to the landscape levels (e.g., Hunsaker et al. 1990). As the literature expanded, the journal *Ecological Indicators* was introduced in 2001, dedicated to the topic. Clearly, there is a plethora of indicators that could be used to characterize ecosystem health, so the issue is not whether indicators of ecosystem health exist but rather identifying the particular sets of indicators that are most efficacious for understanding ecological condition and informing environmental management. We suggest that the specific sets of indicators that should be used logically and directly emerge from the integrated assessment–decision framework, discussed next.

DPSCR₄ FRAMEWORK: NEED FOR A NEW SYNTHESIS FRAMEWORK

Developing a conceptual framework for assessing a system of the scale and complexity of the Gulf of Mexico, with the diversity of audiences that need to be informed,

requires a synthesis of the risk-based and DPSIR approaches, building upon the strengths of each while avoiding their deficiencies. The resulting conceptual framework consists of Drivers–Pressures–Stressors–Condition–Responses elements (DPSCR₄; Figure 1). This framework includes terms that have been used elsewhere in similar contexts, but because there is often inconsistency across the literature in usage of many of these terms, we define each element here to provide the specific meaning of the words as they are used in the DPSCR₄ framework. A few examples of various components of the DPSCR₄ framework applied to the Gulf of Mexico illustrate the construct (Table 1). In actuality, an assessment framework capable of meeting those scale and complexity criteria becomes applicable to virtually any ecosystem and any stressor regime and thus not limited to the Gulf of Mexico.

Drivers

Drivers are the fundamental forces, natural or anthropogenic, that ultimately drive the system. Drivers tend to be large-scale, long-term forces that are not easily controlled or diverted. Examples include demographic drivers (e.g., global population growth or demographic age structure), social drivers (e.g., expansion of human populations into previously undeveloped sensitive habitats), economic drivers (e.g., industrial and energy development), and natural drivers (e.g., the unequal distribution of solar energy across latitudes).




Figure 1. EcoHealth Conceptual Framework (DPSCR₄). DPSCR₄ = Drivers–Pressures–Stressors–Condition–Responses.

Table 1. Examples of DPSCR₄ elements for the Gulf of Mexico

Drivers	Pressures	Stressors	Condition	Responses
Natural and anthropogenic	Human activities and natural processes	Natural and anthropogenic	Assessed on VECs	Societal and ecological
<i>These are the fundamental forces</i>	<i>These are what cause stressors</i>	<i>These are what the ecosystem sees</i>		<i>Reduction, remediation, restoration, and recovery</i>
Economic drivers	Human activities: Resource extraction	Biological stressors	Fish and wildlife VECs	Policies to reduce stressors
			Goal: Sustainable fish and wildlife	Managing drivers and pressures
	<ul style="list-style-type: none"> Commercial fishing Recreational fishing Oil and gas extraction Groundwater usage 	<ul style="list-style-type: none"> Invasive species Overfishing Altered genetics Pathogens Harmful algal blooms 	<ul style="list-style-type: none"> Fisheries populations Avian populations Marine mammals Sea Turtles Endangered species Economic species 	<ul style="list-style-type: none"> Environmental regulations Land-use management Fisheries management Environmental education Conserve special places
Demographic and social drivers	Human activities: Physical	Physical stressors	Habitats	Remediation
			Goal: Restore and sustain habitats	Removing existing stressors
	<ul style="list-style-type: none"> Coastal development Dredging Shoreline structures Transportation Channelization Land-use changes Dams 	<ul style="list-style-type: none"> Habitat alteration Hydrological alteration Changes in salinity Changes in climate Suspended sediments Noise Ocean acidification Hypoxia 	<ul style="list-style-type: none"> Wetlands Mangroves Oyster reefs Seagrasses Coral reefs Barrier islands Freshwater and saltwater marshes 	<ul style="list-style-type: none"> Clean up oil spills Clean up chemical spills Clean up toxic waste sites
Natural drivers	Natural processes	Chemical stressors	Ecological features	Restoration
			Goal: Restore ecosystem health	Restoring ecosystems
	<ul style="list-style-type: none"> Solar energy differences across latitudes Earth's rotation 	<ul style="list-style-type: none"> Nutrient inputs Pesticides Endocrine disruptors Chemical or petroleum spills Trophic dynamics 	<ul style="list-style-type: none"> Connectivity of Gulf with coastal waters Landscape mosaic Biodiversity 	<ul style="list-style-type: none"> Plant seagrasses Restore freshwater flows Increase wetland habitats Remove invasive species
				Ecological recovery
				Ecological processes to return ecological health

DPSCR₄ = Drivers–Pressures–Stressors–Condition–Responses; VEC = valued ecosystem component.

Pressures

Pressures are human activities or natural processes that generate environmental stressors. They also tend to be large scale and long term but often can be highly variable over space and time. Examples of anthropogenic pressures (i.e., human activities) include agriculture, aquaculture, geophysical resource harvesting (e.g., oil exploration and mining), biological resource harvesting (e.g., fishing and forestry), coastal development, marine transport, recreation and tourism, flood control, and the anthropogenic components of global climate change and sea-level rise. Natural processes include ocean dynamic processes (e.g., upwelling and currents), climate processes (e.g., jet stream dynamics, monsoons, and El Niño–Southern Oscillations [ENSO]), sediment dynamics (e.g., subsidence, sedimentation), episodic events (e.g., earthquakes, tsunamis, and hurricanes), and the natural processes components of global climate change and sea-level rise.

Stressors

Stressors are what the ecosystem directly experiences, that is, the physical, chemical, or biological factors that can directly cause an ecological effect. Stressors are the critical point of intersection between the drivers and pressures and the resultant effects on ecological systems; consequently, these are the central cause-and-effect relationships for scientific inquiry and hypothesis testing. Examples of physical stressors include habitat alteration and loss, altered sedimentation and light regimes, altered salinity regimes, drought, hypoxia, and hydrologic alterations. Examples of chemical stressors include oil and chemical spills, altered nutrient inputs, pesticides, and other xenobiotics. Examples of biological stressors include invasive and introduced exotic species, overfishing or overharvesting, pathogens and disease, HABs, and altered genetics. Stressors may secondarily generate other stressors; for example, hydrologic alterations can lead to hypoxia, invasive species, and altered regimes of flooding, sedimentation, turbidity, light, and salinity.

Stressors may involve natural attributes of a system (e.g., the salinity regime of an estuary), which only become stressors when there is a change in the attribute over time or space (e.g., reduced freshwater inflow causing hypersalinity in locations or at times where none previously existed), or they may involve something novel to the ecosystem, such as xenobiotic toxic chemicals or habitat alterations. An environmental stressor may result from one or more pressures or even a mix of natural and anthropogenic pressures. For example, water management that reduces freshwater flows (anthropogenic) and ENSO-induced alterations in precipitation patterns (natural) both can produce a similar stressor (changes in the salinity regime of an estuary). Finally, stressors are system specific, and what is a stressor to 1 ecosystem (e.g., fire in a mangrove forest) may not be a stressor to another ecosystem (e.g., fire in a grassland).

Ecological condition

The state of the ecosystem is its condition or “health.” Because there is an almost unlimited number of specific aspects of an ecosystem that could be used to characterize an ecosystem, a subset of attributes must be identified that are important either ecologically and/or societally, that is, assessment endpoints (USEPA 1998) or VECs (CCME 1996; Harwell et al. 2011). It is advantageous to select a parsimonious set of VECs, with some VECs representative of other similar components of the ecosystem, thereby reducing the number of attributes and causal relationships that need to be characterized to a reasonable and practical set. The set of VECs selected to characterize ecosystem condition should not only focus on endangered or economic species, as is often the case, but also consider ecological scale and hierarchy, and both ecological structure and ecosystem processes. Examples of structural VECs include endangered species, economically important species (e.g., a valuable fisheries population), intertidal or benthic communities, and primary producers. Functional VECs are ecological processes, such as primary productivity, biogeochemical cycling, nutrient dynamics, and trophodynamics. The VECs may also broadly relate to environmental quality, such as water quality, habitat mosaic across the landscape, and biodiversity. Particularly useful for our integrated assessment framework is the subset of VECs that constitutes ecosystem services, including provisioning services (e.g., fish stocks), regulating services (e.g., C storage associated with habitat loss), and cultural services (e.g., bird-watching or other environmentally related recreation and tourism) (UNEP WCMP 2011; Egoh et al. 2012; Hattam et al. 2015).

Finally, in characterizing a VEC (e.g., brown pelican), it may be appropriate to measure the VEC directly (e.g., number of pelicans in a population), but often indicators need to be identified that indirectly reflect on the condition of the VEC. For instance, indicators could include the pelican population age structure, the frequency distribution of eggshell thicknesses, the areal extent and distribution of breeding colonies, or the body burden of PCBs in adult pelicans. Other examples of VECs and associated indicators for the Gulf include the following:

- indicators for VEC water quality: chlorophyll *a*, transparency, total suspended solids;
- indicators for VEC coral community health: coral cover, juvenile recruitment, algal cover, coral composition;
- indicators for VEC seagrass community health: areal extent, seagrass density, nutrient status, community composition; and
- indicators for VEC habitat mosaic: spatial frequency of habitat types and patch-size distributions.

In general, the metrics for each indicator should collectively represent the condition of the VEC at a particular point in time and space.

It is the indicators that will form the foundation of the EcoHealth assessments, including not only indicators that characterize VECs, but also indicators that characterize stressors and pressures, thereby identifying risks to the environment or possible causes for observed effects, plus targets for responses to reduce stressors and improve environmental health. Additionally, the particular levels or trends characterized by the effects indicators can be compared with specific benchmarks, such as historical conditions, desired goals for the particular VEC, or benchmarks between impacted conditions and recovery (Harwell et al. 1996). This comparison allows assignment of qualitative categories of condition, such as degraded, fair, or healthy, or more quantitative ecological health metrics, such as grades, scores, or indices.

Responses

In the original environmental management-based PSR and DPSIR frameworks, response was meant to capture societal feedbacks in response to the ecological impacts, particularly environmental and economic policies and programs intended to prevent, reduce, or mitigate pressures and/or environmental damage (OECD 1993; EEA 1999). In the new framework, we expand responses to include not only such societal responses but also ecological responses, that is, changes in the ecological system. Four types of responses are identified: reduction of stressor sources, remediation of existing stressors, ecological restoration, and ecological recovery.

Reduction. Stressor source reduction consists of societal responses targeted at the management of the drivers and pressures in order to reduce stressors. Examples include policies to reduce greenhouse gas emissions or require more effective wastewater treatment systems. Stressor source-reduction responses may also entail activities such as enhanced environmental educational programs or providing consumers with clearer information on the source and safety of seafood in the markets, among many other examples.

Remediation. Remediation is the set of actions specifically aimed at reduction or elimination of a stressor that has been released into the environment, typically chemical stressors such as toxic wastes or an oil spill; thus, this component reflects the suite of cleanup (i.e., remedial) activities implemented under Natural Resources Damage Assessment (NRDA) regulations (derived from the Comprehensive Environment Response, Compensation and Liability Act [CERCLA 1980]) and the Oil Pollution Act of 1990 (OPA 90 1990) regulations (NOAA 1996a, 2010).

Restoration. Restoration occurs when intervention is made directly into the ecological system in order to undo ecological damage that has been done or to accelerate or enhance the process of ecological recovery, discussed next; it also a component of NRDA regulations (NOAA 1996b). Restoration may entail such actions as removal of invasive

species, reconstruction of wetlands, planting of trees in riparian habitats, adding riffles and pools to a stream, or introduction of an endangered or extirpated species into its former habitat.

Recovery. Recovery, the final “R” in our framework, differs from the others in that it involves natural ecological processes of an ecosystem, once a stressor has been eliminated or reduced below adverse effects levels. Recovery reflects ecological resilience, that is, whether or not and how quickly an ecosystem returns to normal once it is no longer under stress (Holling 1973). Thus, recovery is an internal ecological feedback process, rather than a societal one. An ecosystem has recovered from an incident, such as a chemical or oil spill, once the stressors are gone and all VECs have returned to some baseline condition, given dynamical ecosystem changes and natural variability. Consequently, recovery occurs when there no longer are ecologically significant adverse effects (Gentile and Harwell 1998). The corollary is that recovery cannot fully proceed until the stressors are reduced to below an effects threshold. Where stressors are continuing or periodic, ecological feedbacks may entail permanent changes or even ecological phase shifts in place of recovery. More thorough discussions of ecological recovery are presented in Harwell et al. (2013) and Harwell and Gentile (2014).

Several advantages of the new DPSCR₄ construct underpin ecosystem health assessments: The full sequence of causal relationships is delineated from the ultimate source (fundamental societal or natural drivers) through its manifestation as pressures (human activities and natural processes) and the resulting environmental stressors that the system actually sees, to the effects on ecological condition and the responses that ensue, either through societal actions or natural ecological recovery processes. Second, by taking these relationships from the broad scale down to the specific cause–effect process, an EcoHealth assessment can characterize the system simultaneously from the big-picture policy level to the hypothesis-driven scientific level and back. When nested within a hierarchy of reporting levels, as discussed below (EcoHealth reporting structure), this framework can inform interested audiences at all levels. Similarly, this framework is ideal for aggregation and disaggregation, in which finer scale issues may be explored and illuminated, or in which broader relationships can be more readily perceived. Moreover, this framework can adapt and evolve as more information is gathered and the system becomes better understood, as things change over time or space, or as the scale of the environmental problem varies from small and local to very large scale and multi-dimensional. Thus, the EcoHealth assessment framework can become both responsive to new management needs or questions and useful in identifying uncertainties and new areas of research or monitoring.

The DPSCR₄ framework corrects the 2 serious flaws in the DPSIR framework (i.e., skipping over the critical stressors component and thus missing the causal part of cause-and-effects relationship, and the conflation of state and impacts).

Finally, the DPSCR₄ conceptual framework provides the basis and rationale for identifying the specific sets of indicators in EcoHealth assessments for pressures, stressors, and condition, the particular suite of attributes desired for each indicator and VEC, and insights into the management or societal actions that could be implemented to achieve ecological health goals; consequently, the DPSCR₄ framework provides guidance both for ecosystem health assessment and for environmental decision making. In essence, the DPSCR₄ framework functions as a systematic and comprehensive set of sieves, partitioning and filtering information about the coupled human–ecosystem of concern, from which emerges the specific set of indicators to comprise an ecosystem health report card or, as discussed next, the complete set of EcoHealth metrics (bottom of Figure 1).

EcoHealth reporting structure

The integrated ecosystem assessment framework is further structured to inform a diversity of audiences with differing concerns and levels of scientific understanding. This is accomplished by overlaying onto the DPSCR₄ framework a structural hierarchy that emphasizes tier-relevant components and indicators (Figure 2) appropriate to the differing types of audiences to be informed by an ecosystem health assessment. The top level is the target of the original PSR and DPSIR frameworks, focused on the overall condition of the environment, the broad pressures that influence it, and the societal responses that ensue. It requires very few indicators of health and thus constitutes the greatest degree of aggregation into the most simple-to-understand synthesis metrics and formats, that is, a report card.

The next lower level is the realm of people who make or influence environmental decisions and policy. This tier emphasizes impacts from pressures on the environment and specific societal responses to mitigate impacts by managing pressures. This level requires more information because the audience tends to be more engaged in the issues of concern. Below that is the level of hands-on environmental managers, for example, managing a park or conservation lands. These individuals need to understand a diversity of environmental issues relevant to their specific location or ecosystem types. Consequently, this tier focuses on specific stressors, their impacts on particular ecosystems of concern,

and remediation and restoration activities that might be implemented to achieve management goals.

At the base of the hierarchy is the scientific community whose hypothesis-driven focus is on environmental stressors, their effects on ecological condition, and whether the effects constitute adverse health compared to baseline or benchmark conditions; remediation and restoration activities to improve the health of the environment; and the ecological processes underlying ecosystem recovery and determining when recovery has been attained. Indicators at this tier are numerous and aggregation is minimal, consistent with the many hypotheses concerning stress–effects relationships in ecosystems. These indicators are the ultimate foundation of the EcoHealth assessments, that is, specific qualitative or quantitative metrics that reflect the relevant characteristics of each VEC and of each pressure and stressor over time and space. The utility of each indicator depends on fidelity to condition, data availability, ability to interpret and explain results, and spatial and temporal applicability (Kelly and Harwell 1989, 1990; Dale and Beyeler 2001). Development of databases and monitoring for each indicator, including establishment of reference or benchmark conditions (Harwell et al. 1996; Harwell, Gentile et al. 1999; Harwell, Myers et al. 1999), can provide the foundation for understanding the dynamics of each VEC, its trajectory over time and space, and its health or recovery status.

The hierarchical reporting structure presents a dynamic framework for aggregating information into more integrative indicators at higher levels and for channeling specific information requests from higher tiers down to the appropriate level. As information is acquired by scientific investigations or through environmental monitoring, updated or new indicators can be provided to the tiers above. Concomitantly, information needs identified at higher levels can guide the scientific investigations performed, inform the allocation of resources to reduce important uncertainties, or encourage development of new integrative metrics. Thus, the DPSCR₄ hierarchy provides the template for this 2-way information exchange to occur and may lead to more efficacious acquisition and utilization of research and monitoring data. Similarly, it avoids the limitation of the strictly stress-effects–based assessment framework because its hierarchical structure and aggregation or disaggregation processes allow it to better accommodate multiple stressors.




Figure 2. EcoHealth metrics hierarchical reporting structure.

Constructing a Gulf EcoHealth report card

A long-term objective is to apply the DPSCR₄ framework to assess the ecosystem health of the Gulf of Mexico. The strategy is to disaggregate the Gulf into manageable regions, subregions, or other reporting units, and then sequentially follow the DPSCR₄ framework to develop the Gulf EcoHealth report card specific to that unit. The Gulf can be partitioned into several regional-scale subunits based on geographical, ecological, and/or political boundaries. Within each region are delineated specific habitat types of concern, such as seagrass or salt marsh communities, and within each habitat are identified specific sets of VECs and associated indicators. Moreover, crosscutting ecological components, such as migratory birds and marine mammals, are very relevant to ecosystem health and thus essential components of EcoHealth assessments. When completed, these component assessments would be reaggregated across spatial and ecological scales, essential for characterizing the health of such a large and complex ecosystem as the Gulf of Mexico.

We developed a stepwise process for constructing the EcoHealth report card for any particular area:

- 1) Develop the conceptual ecosystem model. This conceptual modeling process should involve scientists, managers, and stakeholders familiar with the systems of concern to ensure that ecosystem elements are adequately identified and long-term sustainability goals are appropriately defined. Typically, this is done in a workshop setting, in which the region or subregion is partitioned into its constituent ecological habitats, and the VECs of each habitat are identified, along with the drivers, pressures, and stressors impinging upon them. Next, a series of matrices are constructed linking the drivers or pressures and their resulting stressors with the VECs that would be affected, including assigning a relative strength of each stressor–effect relationship. An example of such a matrix is illustrated in the pilot study on coastal Texas rookery islands presented in the next section. By having knowledgeable workshop participants make such judgments on the interactions of every stressor and every VEC for each habitat in the ecosystem of concern, a substantial amount of information is captured about how the ecosystem responds to natural and anthropogenic stressors and the pressures and drivers that cause them. From these completed matrices a set of habitat-specific, risk-based graphical conceptual ecosystem models (CEMs) can be constructed; collectively, the CEMs for a region should reflect the connectivity among all the ecosystem components. In the format we use for these conceptual models (Supplemental Data Figure 1), the top tier (shown as rectangles) is pressures, in this case human activities that impinge on the Mission-Aransas Reserve, Texas, USA landscape. The next tier (ovals) is the environmental stressors that result from the pressures to which they are linked in the graphic, with thicker lines representing stronger linkages. At the bottom tier (hexagons) are the VECs, identified here for the landscape-level attributes of the Reserve, showing the weighted linkages between specific stressors and effects on specific VECs. For additional examples of this risk-based class of CEMs, see Cormier et al. (2000), Gentile et al. (2001), and Ogden, Davis, Barnes et al. (2005), and Ogden, Davis, Jacobs et al. (2005).
- 2) Select indicators. For each conceptual model derived from the DPSCR₄ framework, indicators are identified for spatially explicit reporting on each important pressure, stressor, and VEC. Selected indicators should be data driven, reliably measurable, and/or based on integrative techniques. Collectively, the goal is for a parsimonious set of indicators that captures the information needed to characterize and evaluate ecosystem health, reflecting current status and future trends for pressures or stressors and for ecological condition. Indicators should be chosen with consideration of their use within the EcoHealth assessments (Table 2) (Kelly and Harwell 1989, 1990).
- 3) Define goals, benchmarks, and thresholds for assessment. Goals are defined here as the desired condition for the particular ecosystem or ecosystem component, often identified in the context of ecological sustainability. Benchmarks are defined here as milestones along the way from the current condition toward the desired sustainable state (Harwell, Myers et al. 1999). Thresholds may be identified that mark particular levels of health, often useful for communicating ecosystem condition. A quantitative or qualitative metric that defines a desired condition or goal for each indicator should be established, allowing indicator metrics to be assessed and reported in the EcoHealth assessments. Goals and benchmarks can be set in several ways, including using established regulatory metrics (e.g., numerical ambient water quality criteria; Stephan et al. 1985), identifying biologically or ecologically relevant data values from the literature (e.g., defining hypoxia to be $\leq 2.0 \text{ mgL}^{-1}$ dissolved O; Rabalais et al. 2002), comparisons to historical conditions prior to major impacts (e.g., assessing areal coverage of seagrass communities in the northern Gulf; Carter et al. 2011), or measurements of benchmarks that have been achieved in similar ecosystems elsewhere.
- 4) Characterize results. Indicator values are evaluated against specific goals, benchmarks, or thresholds. These may be standardized into assigned condition categories, and values for individual indicators may be integrated to produce an overall index or other metric for the VEC, pressure, or stressor. These may be spatially integrated to characterize a subregion or region of concern, and these in turn may be further integrated with other subregion or region results using an area-weighting approach.
- 5) Communicate results. The communication of results is the ultimate goal of the EcoHealth assessments; it should be multifaceted and transparent, structured hierarchically for different audiences (Figure 2). Each EcoHealth assessment document should be a graphics-rich, synthesis document that aggregates results to

Table 2. Purposes and criteria for selecting indicators

Purpose of indicators	Criteria for selecting indicators
Intrinsic importance—key: indicator is the endpoint <ul style="list-style-type: none"> • Examples: economically important species; endangered species 	Signal-to-noise ratio <ul style="list-style-type: none"> • Sensitivity to stressor • Intrinsic stochasticity
Early-warning indicators—key: rapid indication of effects <ul style="list-style-type: none"> • Screening tool • Quick response time • Low signal-to-noise ratio, low discrimination • Accept false positives 	Rapid response <ul style="list-style-type: none"> • Early exposure • Quick dynamics (e.g., short life span)
Diagnostic indicators—key: reliability in predicting effects <ul style="list-style-type: none"> • High stressor specificity • High signal-to-noise ratio • Minimize false positives 	Reliability or specificity of response <ul style="list-style-type: none"> • Available field protocols • Preexisting database • Low-cost tools
Process or functional indicators—key: process in the indicator <ul style="list-style-type: none"> • Monitoring other than biota (e.g., decomposition rates) 	Ease or economy of monitoring <ul style="list-style-type: none"> • Answers the “so what” question
	Feedback to managers

Modified from Kelly and Harwell (1990), with permission from Springer Nature. © 1990.

create an easily understandable message about the overall health of the ecosystem. Underlying information, source documents, and linkages to data sources are important to providing transparency of process and accessibility to information appropriate for managers, decision makers, program managers, and scientists. Assessment results may be communicated on an annual and/or multiyear cycle, similar to the series of Chesapeake Bay Report Cards and Everglades System Status reports (e.g., IAN 2007, 2013; ecoreportcard.org; evergladesrestoration.gov).

TESTING THE CONCEPTUAL FRAMEWORK: TEXAS COASTAL ECOSYSTEMS REPORT CARD AND MISSION-ARANSAS RESERVE ROOKERY ISLAND PILOT PROJECT

Because an EcoHealth Metrics for the Gulf of Mexico is so complex and diverse, we conducted a proof-of-concept evaluation of the DPSCR₄ conceptual framework and its implementation for assessments through constructing a prototype Texas coastal ecosystems report card. Coastal Texas and its watersheds provided an excellent pilot project for the entire Gulf because of the diversity and complexity of its ecosystems, human communities, and associated environmental pressures and stressors. In the present pilot study we followed the 5 steps listed in the preceding section, focusing on a manageable subset of Texas coastal ecosystems and crosscutting ecological components, specifically seagrass meadows, oyster reefs, recreational and commercial fisheries, and resident and migratory coastal birds. The resulting prototype Texas Coastal

Ecosystems Report Card (McKinney et al. 2018a, 2018b) provides details on the pressures, stressors, VECs, ecosystem services, stress-effects matrices, CEMs, assessment data, scoring methodologies, and results for each system type evaluated; a summary of the fisheries and avian report cards are presented in the Supplemental Data.

This Texas pilot study demonstrated that the framework is ideal in providing a systematic and comprehensive methodology for understanding the coupled human-ecological systems and for identifying the elements of a comprehensive ecosystem health report card. Unfortunately, the available data for ecosystems assessment were limited to only a few selected attributes for some VECs of some systems, and other than water quality information, which has been collected at length by various state and local agencies, only minimal information exists on most pressure and stressor indicators for any ecosystem. As a result, we could not construct the Texas report card called for by the DPSCR₄ framework. From existing data we could derive the status and trends of the health of selected fish and avian species (Figures 3 and 4, respectively). However, the identical report card on these 2 ecosystem components could have resulted had we simply used existing protocols for report cards, such as the Chesapeake Bay and Mississippi River systems (e.g., IAN 2013; americaswatershed.org). Consequently, the DPSCR₄ conceptual framework provided no added value for conducting ecosystem health assessments when using existing data because of the paucity of data. On the other hand, the DPSCR₄ framework was very useful in identifying the elements of the environmental monitoring program that would be necessary for comprehensively assessing ecosystem health of the Gulf of Mexico and progress toward sustainability goals; that is, its essential utility is prospective, guiding future data acquisition and analyses. This is especially important given the extensive financial resources that are becoming available for Gulf of Mexico environmental research and monitoring resultant from the settlement of the Deepwater Horizon legal (e.g., Clean Water Act penalties) and legislative processes (e.g., the RESTORE Act 2012 and RESTORE Act 2015; see <https://www.restorethegulf.gov/history/about-restore-act>).

The second proof-of-concept pilot study was designed to examine the DPSCR₄ assessment and decision framework as applied to a specific management issue of the Mission-Aransas Reserve (<https://missionaransas.org>; Figure 5). Mission-Aransas is one of 29 National Estuarine Research Reserves around the country, each a federal and state partnership for conducting research, education, and stewardship programs. The Mission-Aransas Reserve, which is a partnership between National Oceanic and Atmospheric Administration (NOAA) and the University of Texas Marine Science Institute, has as its primary goals to improve knowledge of the ecosystem structure and function of Texas coastal ecosystems, promote understanding of the Texas coastal ecosystems by diverse audiences, and promote public appreciation and support for stewardship of coastal resources (Mission Aransas Reserve 2019).




Figure 3. Status and trends for selected fish and invertebrate species.

The present pilot study was designed to use the components of the DPSCR₄ conceptual framework to characterize the critical linkages between the ecological and societal systems, identify specific key indicators for assessing ecological health and ecosystem services, identify management alternatives for assessment, and apply ecosystem-based management tools to analyze the options with respect to achieving management goals. We asked the managers and scientists from the Reserve to identify an important environmental management issue that might benefit from applying our integrated assessment framework for guidance. The management goal selected for the pilot study was to create or expand rookery islands in order to

enhance nesting, breeding, and foraging habitat for both resident and migratory coastal birds. Colonial waterbirds function both as a critical ecosystem services component and as an important ecological functional component of the Mission-Aransas ecosystem.

The management focus on enhancement of rookery island habitat results from the risk to the long-term sustainability of resident and migratory coastal birds by the historical and ongoing degradation or loss of critical habitat in the coastal areas of the Mission-Aransas Reserve and the associated Coastal Bend Bays & Estuaries Program (CBBEP; <http://www.cbbep.org>). Analysis of data from 1973 to 2008 on 14 colonial waterbird species that




Figure 4. Status and trends for selected breeding and wintering bird species.

use the Texas Coastal Bend area shows that the populations of several species are decreasing, driven primarily by a decrease in available waterbird island nesting habitat, resulting from sea-level rise and storm-surge erosion, subsidence, increasing human pressure and habitat loss adjacent to both nesting and feeding areas, and a scarcity of adequate nesting substrate. Consequently, Reserve and CBBEP managers and scientists identified suitable nesting habitat to be the major limiting factor in the long-term sustainability of waterbird species in the Coastal Bend area (CBBEP 2010). Colonial-nesting waterbirds require islands for breeding that provide suitable nesting habitat (e.g., shrubs for wading birds, bare ground for

terns) free from predators and disturbance sources and relatively close to feeding areas. Unlike barrier islands in the region, the rookery islands are typically smaller and consequently less affected by predators (CBBEP 2010), and thus became the focus of the pilot study.

The pilot study approach was to apply a structured decision-support system to the DPSCR₄ framework and use geospatial ecosystem-based management tools and scenario-consequence analyses to assess the management options. We systematically characterized each of the DPSCR₄ components for rookery islands (Table 3), which suggested stressors to be minimized and attributes to be optimized in rookery island site identification and selection.




Figure 5. Map of the Mission-Aransas National Estuarine Research Reserve. NERR = National Estuarine Research Reserve; SWMP = System-Wide Monitoring Program.

We began with the development of conceptual ecosystem models for the rookery islands by examining the stressor–effects relationships as shown in the causal matrices for physical, chemical, biological, and climate-change stressors (Table 4). Each entry in the matrix reflects how large an effect, if any, on each VEC would occur by the presence of each stressor at current or plausibly anticipated levels of intensity, duration, and extent. For example, the stressor changes in the hydrological regime was expected to have a high effect on rookery island areal extent and/or location, structural complexity, successional patterns, and habitats for resident and migratory birds, whereas the stressor marine debris was not expected to have any effects on areal extent and only low-level effects on the other structural attributes. By systematically capturing the consensus expert opinion on the strength of stressor–effects relationships for all stressors and all VECs of a system, the risk regime of that system is essentially characterized. The complete rookery island CEMs are shown in Supplemental Data Figure 2.

The key factors in rookery island habitat emerge from the application of the DPSCR₄ framework to the Mission-Aransas environmental issues and the development of the details of Table 4, including 1) the Reserve management's emphasis on ecosystem services and well-being

highlighted the particular value of rookery islands to support bird-watching and associated activities in the Texas Coastal Bend region; 2) the structural attributes of rookery islands directly affect their functional attributes; 3) a number of important stressors can adversely affect the spatial extent and habitat quality of rookery islands, which are critical to sustainability of the coastal bird populations; 4) specific site characteristics affect the potential of rookery islands to achieve desired attributes (e.g., distance from major land mass in order to preclude invasion by predators that often seriously inhibit or prohibit nesting birds); and 5) the rookery island site-selection process was guided by the DPSCR₄ framework toward reducing those specific stressors and optimizing those desired structural and functional attributes.

Two management scenarios were explored in the Mission-Aransas pilot study: 1) enhancing existing rookery islands in order to improve the availability of habitats for nesting and/or migratory birds and 2) identifying the optimal location and configuration for new artificial rookery islands for single- or multipurpose use (e.g., adding oyster reefs or recreational fishing habitat). The spatially explicit Texas Colonial Waterbird Society data for 1970 to 2015 were acquired for specific locations of islands or island

Table 3. The DPSCR₄ framework populated for Mission-Aransas rookery islands

Stressors	Condition attributes	Ecosystem services	Well-being attributes
Physical stressors	Structural attributes	Bird watching	Recreation
Changes in salinity regime	Areal extent	Recreation	Economic
Changes in precipitation	Habitat diversity	Navigation	Cultural
Erosion	Structural complexity	Habitat value	Health
Sea-level rise	Successional patterns	Biodiversity	
Inundation	Breeding resident birds		
Storms	Winter-migratory birds		
Noise	Functional attributes		
Chemical stressors	Colonial waterbird breeding habitat		
Nutrients	Waterbird nonbreeding habitat		
Petroleum releases	Whooping crane habitat		
Pesticides and herbicides	Marsh habitat		
Biological stressors	Seagrass habitat		
Food availability	Fish habitat		
Predation	Invertebrate habitat		
Harmful algal blooms	Oyster habitat		
Human presence	Erosion protection		

DPSCR₄ = Drivers–Pressures–Stressors–Condition–Responses.

clusters within the Reserve and surrounding area (TCWS 2019). Additional geospatial data were acquired for important landscape variables (e.g., distance and direction from land masses and sources of predators), hydrological attributes (e.g., bathymetry), and biological attributes (e.g., benthic cover, presence of existing rookery site) that could be used to identify candidate sites for rookery island creation or enhancement.

These geospatial data were layered into the geographical information system (GIS) software extension package, NatureServe Vista (NatureServe 2019), to develop an index that characterizes the Mission-Aransas Estuary system and adjacent estuary systems vis-à-vis their suitability for rookery island development. Suitability factors related to rookery island creation and/or enhancement (e.g., water depth, distance from shoreline, benthic habitat cover) were determined with input from local experts and were characterized spatially across the landscape on the basis of their assigned suitability characteristics (Table 5). The suitability index was calculated from the weighted values of each relevant suitability factor. NatureServe Vista was then used to determine where the most suitability factors for rookery island creation or enhancement were found together (for details, see Stanzel 2017).

Suitability scores were calculated for the entire study area. The highest priority sites (i.e., specific areas where the most suitability factors overlapped) were identified using the Site Explorer tool within NatureServe Vista. If the high-suitability grid cells were located adjacent to another high-suitability grid cell with the same score, multiple cells were aggregated, resulting in 30 high-suitability candidate sites. To allow for better visualization to distinguish existing rookery island sites that are suitable for enhancement versus high-priority sites for creation of a new multipurpose rookery island, each of the high-suitability candidate sites was identified using a single point in the grid.

Each of the 30 high-suitability sites were further characterized using the following information: 1) suitability score, 2) size (number of acres), 3) location (latitude/longitude), 4) average water depth, 5) benthic habitat type, 6) average distance from shoreline, 7) average direction from shoreline, 8) water with depth of 4 feet located within 800 feet of site, and 9) presence of an existing rookery. These characteristics proved to be helpful to managers as they prioritized within the list to determine which sites should be further explored for their potential for restoration or enhancement. For example, “water with depth of 4 feet within 800 feet” suggests which sites may be more accessible by barges carrying material for island creation or restoration, and characterization of the nearby habitat can help determine the likelihood that mitigation for damage to seagrass or oysters caused by island construction would be required.

In a final workshop, scientists and managers from the Harte Research Institute, the Mission-Aransas Reserve, CBBEP, and other state and federal agencies used the index results and the suitability criteria to select priority sites for additional investigation (Figure 6). For enhancing existing rookery island sites, Deadman’s Reef and Third Chain of Islands were assigned highest priority, and Redfish Bay and Big Bayou Spoil were identified as worthy of further exploration. For creating a new rookery island, top priority was assigned to Seadrift in San Antonio Bay, and the sites deemed worthy of further exploration were Falcon Point, Second Chain of Islands, and Cape Carlos.

Unfortunately, shortly after these analyses were completed in August 2017, Hurricane Harvey passed directly over the study area. A post hurricane reconnaissance survey of rookery islands in the area by staff from CBBEP found widespread destruction, including near-complete loss of several islands, severe habitat degradation on nearly all the islands, and loss of almost all protective signage, along with

Table 5. Summary of conservation elements and suitability criteria used in the rookery Island pilot study

Conservation element	Rationale	Suitability	Weighting
Water depth	Water depth affects the cost or feasibility of rookery island creation: As the water gets deeper, it becomes cost prohibitive to create above-water habitat.	Water depth \leq 2 ft = 1.0 Water depth 2–4 ft = 0.7 Water depth 4–6 ft = 0.2 Water depth \geq 6 ft = 0.0	1.0
Distance from shoreline	Distance of the rookery from mainland shorelines affects the ability of predators to reach the island: The farther an island is from other shorelines, the less likely it is to be invaded by predators.	Distance of rookery from shoreline < 0.5 mi = 0.0 Distance of rookery from shoreline \geq 0.5 mi = 1.0	1.0
Direction from shoreline	Because of locally prevailing southeasterly winds, the direction from mainline shorelines affects the ability of predators to sense a rookery.	Southeast = 1.0 East = 0.8 South = 0.5 All other directions = 0.0	0.5
Habitat	The presence of seagrass and oysters will result in the need for habitat mitigation, which increases project costs. Locating islands in areas where mitigation potential is lower is a desired attribute.	Seagrass present = 0.0 Oyster present = 0.0 Unconsolidated bottom = 1.0 Unknown = 0.8	1.0
Location of known rookery islands	Location of active and inactive rookeries could increase the potential for restoration and /or enhancement.	Active rookery = 1.0 Inactive rookery = 0.5	1.0

deposition of large debris across former nesting habitats (Newstead and Fitzsimmons 2017). The specific sites that we had identified as priority sites were in the hardest-hit areas and were severely damaged or destroyed. The scientists implemented a rapid response to do as much habitat restoration as possible during the subsequent several months before the nesting season was to begin the following spring. This storm not only illustrated the high vulnerability of those sites but also redoubled the need for rookery island restoration as a critical component in recovery and sustainability of Texas colonial waterbirds. Fortunately, our pilot study results were immediately available to assist in the restoration process for recovery from the hurricane, and the tool is ready for longer-term application to implement rookery island nesting habitat restoration.

NEXT STEPS: EXPANSION TO THE GULF OF MEXICO AND BEYOND

The next steps regarding this effort to develop a continuing ecosystem health reporting system for the Gulf of Mexico is expected to be in 2 parts. The initial Texas prototype report card, as summarized here, will be updated with more detailed data and analyses and extended to include an independent section on water quality, which was identified as an important stressor in each of the prototype assessments. This will be issued as a Texas coastal ecosystems report card intended for broad public consumption

and continuing thereafter as a sustainable, ongoing report of ecosystem health for the Texas coast.

Secondly, a goal of the authors is to develop a sustainable report card for the whole Gulf of Mexico. The Texas prototype and its successor report card will be the model for a broader Gulf of Mexico Report Card. The means and methods to do so were an important outcome of the Texas exercise and “proof of concept” that will lay the foundation for that effort. The initial exercise of expanding to the Gulf of Mexico will require an expert workshop to partition the Gulf into units that have both ecosystem and resource management meaning. A follow-up task will be to agree on appropriate indicators that match the scale of the Gulf as a large marine ecosystem but that also have resource management meaning. Again, the prototype provides guidance in accomplishing this task and the subsequent analytical process to develop assessments that can be sustained. An important challenge will be to develop a reporting format and style that are appropriate to the expected and diverse users in the Gulf of Mexico.

In addition to expansion of the EcoHealth assessment framework, we also envision adoption of the DPSCR₄ framework in a much broader context as a generalized conceptual model of the coupled human–ecological system, with applicability across a full suite of ecological systems, natural and anthropogenic pressures and stressors, and management issues. Much as the US Environmental Protection Agency ERA framework and guidelines (USEPA 1992, 1998) have been




Figure 6. Priority sites selected for creating or enhancing rookery islands.

applied to a wide diversity of ecological systems and risk management problems (e.g., OSTP 1999; Suter 2007), we assert that the DPSCR₄ framework provides a systematic and comprehensive guidance to understanding environmental issues and managing for desired outcomes.

As one example of the applicability of the DPSCR₄ conceptual framework, the risk assessment guidelines emphasize the importance of making informed judgments regarding the purpose, scope, and technical approaches early when planning a risk assessment. During the initial planning of an ERA, the manager and risk assessor collaborate to determine who or what is at risk; what are the environmental hazards and stressors of concern, what are the sources of the hazards and stressors, and what are the potential routes of exposure and valued ecological components that are potentially at risk and the focus of the assessment. Thus, the DPSCR₄ conceptual framework provides a systematic and comprehensive construct for the problem formulation phase of the risk assessment, identifying the full spectrum of human and natural pressures, the hazards and stressors that the pressures create, the VECs that potentially are exposed and at risk, and the ecosystem and societal services they provide, including their links to human well-being. This process unfolds as the risk assessor gathers the information to populate the DPSCR₄ (e.g., Table 3).

The populated DPSCR₄ conceptual framework is valuable as an organizing framework for the problem formulation phase of an ERA, thereby providing a comprehensive and systematic picture of the risk landscape for a particular problem. It is also an invaluable tool for conducting a relative-risk

ranking of the potential risks, that is, a process for identifying the most important risk hypotheses needing further examination in the analysis phase of an ERA process. As potentially high-priority risks are identified, the risk assessor can prepare a series of conceptual ecological models to illustrate the exposure pathways linking the pressure to stressors to the structural and functional ecological and ecosystem services endpoints. By identifying the high-priority risks, the DPSCR₄ conceptual framework can facilitate reducing the dimensionality of the risk landscape, allowing the decision makers and risk assessors to focus on the parsimonious set of risk hypotheses that will then become the subject of detailed examination as part of the ERA's analysis phase, which seeks to understand causal relationships and reduce associated uncertainties (USEPA 1992, 1998).

Acknowledgment—The work contained in this manuscript is adopted from research work conducted by the first 6 co-authors and reported in a white paper publicly available online at <https://tinyurl.com/y7vfdlax>. That white paper presented an earlier version of the framework that was provided to workshop participants as a part of the framework development process. The feedback received from the scientists and practitioners at the workshop was then used to modify and finalize the conceptual framework described in the present manuscript. The Harte Research Institute's Gulf EcoHealth Metrics Initiative has been supported in part by the Harte Research Support Foundation, Texas OneGulf Center of Excellence, and the NOAA RESTORE Program. Additionally, the seminal workshop to develop the EcoHealth Metrics framework and initiate the

Texas Pilot Project was supported by the Office of the Governor of the State of Texas. The extension of the framework to include ecosystems services and the Mission-Aransas pilot project were funded by a competitive research grant from the NOAA RESTORE Science Program, NA15NOS4510288. We also acknowledge partial support from the Walton Family Foundation and the Harte Research Support Foundation for assistance in the development of the initial conceptualization of the EcoHealth assessment framework.

We thank the participants (listed in the Supplemental Data) of the 4 workshops convened at the Harte Research Institute for Gulf of Mexico Studies (HRI) in support of this study, including the initial report card conceptual workshop in 2011; the 2 workshops in 2016 to develop and refine the integrated assessment/decision framework, initiate the prototype report card, and initiate the Mission-Aransas pilot study; and the final NOAA RESTORE workshop in which the suitability index-defined candidate rookery island sites were analyzed and selected. We greatly appreciate the contributions of all the workshop participants; a list is provided in the Supplemental Data.

We particularly thank Chris Onuf (retired), who led the prototype report card development for seagrass communities, Jennifer Pollack (Department of Life Sciences, Texas A&M University–Corpus Christi), who led the prototype report card development for oysters, and Beau Hardegree (USFWS), Owen Fitzsimmons (CBBEP), and David Newstead (CBBEP), who provided essential avian and geospatial physical data and expert analyses for the rookery island site suitability/site selection process.

Disclaimer—The authors declare no conflicts of interest.

Data Accessibility—The only newly created data in this study were the stress–effects relationships for the stressors and VECs of the rookery islands of Mission-Aransas Reserve, which were based on expert judgment and are fully reported in Table 4.

The data used in the prototype Texas coastal fisheries report card were derived from publicly available status reports on fish stocks, as detailed in Stunz et al. (2018). Co-author GW Stunz is the contact person for readers requesting additional information (Greg.Stunz@tamucc.edu). The data used in the prototype Texas coastal birds report card were derived from publicly available bird count data, including the Audubon Christmas Bird Count and the North American Breeding Bird Survey Data (<https://www.pwrc.usgs.gov/bbs/>); details are provided in Withers (2018). Co-author K Withers is the contact person for readers requesting additional information (Kim.Withers@tamucc.edu). The complete prototype Texas Report Card and supporting technical support documents are available online at <https://www.harresearchinstitute.org/project/gulf-mexico-report-card-prototype-texas>; the technical support documents provide the details on the pressures, stressors, VECs, ecosystem services, stress-effects matrices, CEMs, assessment data, scoring methodologies, and results for each system type evaluated.

The Mission-Aransas rookery island site-selection suitability factors were derived from publicly available geospatial data based on the Texas Colonial Waterbird Society data for 1970 to 2015 on 14 colonial waterbird species for the Texas Coastal Bend area; additional existing publicly available geospatial data were used for important landscape variables (e.g., distance and direction from land masses), hydrological attributes (e.g., bathymetry), and biological attributes (e.g., benthic cover) for the study area. These data were entered and the analyses were conducted by co-author KM Stanzel using the NatureServe Vista GIS software package; details of the data sources and suitability factors calculations are presented in the text and in Stanzel (2017). Stanzel is the contact person for readers requesting additional information (kstanzel@cbbep.org).

The conceptual ecosystem model for human activities affecting the Mission-Aransas landscape-level ecosystem, cited in the text as Supplemental Data Figure 1, is available through figshare; the complete set of conceptual ecosystem models for the rookery island ecosystems, cited in the text as Supplemental Data Figure 2, is available through figshare.

SUPPLEMENTAL DATA

Supplemental Data Coastal Texas Report Card is a summary description of the prototype Texas coastal report card for the fisheries and avian components. This is the section that was removed from the initial manuscript in response to a review comment, both to shorten the manuscript and to reflect the absence of adequate data to fully create the Texas report card.

Supplemental Data Figure 1 presents an example conceptual ecosystem model for the Mission-Aransas National Estuarine Research Reserve to illustrate the types of conceptual ecosystem models described in the manuscript.

Supplemental Data Figure 2 is the complete set of conceptual ecosystem models for the rookery islands of the Mission-Aransas Reserve and Texas Coastal Bend regions.

Supplement Data Workshop Participants provides the full list of participants for all workshops.

We thank the participants of the four workshops convened at the Harte Research Institute for Gulf of Mexico Studies (HRI) in support of this study, including the initial report card conceptual workshop in 2011; the 2 workshops in 2016 to develop and refine the integrated assessment/decision framework, initiate the prototype report card, and initiate the Mission-Aransas pilot study; and the final NOAA RESTORE workshop in which the suitability index-defined candidate rookery island sites were analyzed and selected.

Collectively, these workshop participants included: Becky Allee (NOAA RESTORE Program Office); Ray Allen (Coastal Bend Bays & Estuaries Program, CBBEP); Porfirio Alvarez (Consortium of Marine Research Institutions of the Gulf of Mexico); Patrick Biber (Gulf Coast Research Laboratory, University of Southern Mississippi); Wylie Barrow (USGS); Melissa Brewer (TAMUCC-HRI); Joanna Burger (Department of

Ecology, Evolution, and Natural Resources, Rutgers University); Ed Buskey (Mission-Aransas Reserve); Cristina Carollo (TAMUCC-HRI); Billy Causey (National Marine Sanctuaries Program, NOAA); Just Cebrian (University of South Alabama); Mark Dumesnil (The Nature Conservancy); Ken Dunton (University of Texas Marine Science Institute); Owen Fitzsimmons (CBBEP); Steve Fletcher (United Nations Environment Programme); Mark Fisher (Texas Parks & Wildlife Department); Jeff Francis (TAMUCC-HRI); Tom Frazer (School of Forest Resources & Conservation, University of Florida); James Gibeau (TAMUCC-HRI); Holly Greening (Tampa Bay NEP); Liz Gomez (NOAA); Larry Handley (USGS); Beau Hardegree (USFWS); David Hicks (University of Texas-Rio Grande Valley); Chris Kelble (NOAA); Chris Mace (Texas Parks & Wildlife Department); Gary Matlock (NOAA); Richard McLaughlin (TAMUCC-HRI); Paul Montagna (TAMUCC-HRI); David Newstead (CBBEP); Amy Nunez (Texas General Land Office); John Ogden (National Audubon Society, deceased); Jennifer Pollack (Department of Life Sciences, Texas A&M University-Corpus Christi); Warren Pulich (Meadows Center for Water and the Environment, Texas State University-San Marcos); Victoria Ramenzoni (TAMUCC-HRI); John Rappole (Conservation Biology Institute, Smithsonian Institution); Denise Reed (The Water Institute, currently Laboratory of Coastal Restoration Science, University of New Orleans); Mike Reiter (Department of Integrated Environmental Science, Bethune-Cookman University); Chris Robbins (Ocean Conservancy); Lance Robinson (Texas Parks & Wildlife Department); Marc Russell (USEPA); Joe Saenz (USFWS); John Schalles (Creighton University); Dana Sjostrom (Mission-Aransas Reserve); Elizabeth Smith (International Crane Foundation); Kiersten Stanzel (Green Wing Environmental, LLC, currently CBBEP); Greg Stunz (TAMUCC-HRI); Katie Swanson (Mission-Aransas Reserve); John Tirpak (USFWS); Jim Tolan (Texas Parks & Wildlife Department); Jace Tunnell (Mission-Aransas Reserve); Wes Tunnell (TAMUCC-HRI, deceased); Aswani Voley (College of Arts & Sciences, University of North Carolina-Wilmington); Hongqing Wang (National Wetlands Research Center, USGS); David Weinstein (Department of Natural Resources, Cornell University); Keith Westlake (USFWS); Mike Wetz (Department of Life Sciences, Texas A&M University-Corpus Christi); Kim Withers (Department of Life Sciences, Texas A&M University-Corpus Christi); Chris Wood (eBird Project, Cornell Lab of Ornithology); Mark Woodrey (Grand Bay NERR); David Yoskowitz (TAMUCC-HRI).

REFERENCES

America's Watershed Initiative. 2015. Mississippi River Watershed report card. America's Watershed Initiative, St. Louis (MO). 8 p. [accessed 2019 Mar 11]. <http://americaswatershed.org/reportcard/>

Australian and Queensland Governments. 2010. Great Barrier Reef first report card 2009 baseline reef water quality protection plan. Reef Water Quality Protection Plan Secretariat, Queensland Department of Environment and Heritage Protection, Brisbane City, Queensland, Australia. 4 p. [accessed 2019 Mar 11]. <http://www.cbbep.org>, http://ian.umces.edu/press/assessment/great_barrier_reef/

Barrett GW, Van Dyne GM, Odum EP. 1976. Stress ecology. *BioScience* 26:192–194.

Brandt LA, Browder JA, Cherkiss M, Frederick P, Gaiser E, Gawlik D, Geiger S, Kelble, Kelly CS, Kline J, Kotun K et al. 2016. System-wide indicators for Everglades restoration. 2016. Technical report. Davie (FL): South Florida Ecosystem Restoration Task Force. 98 p. [accessed 2019 Mar 19]. https://www.evergladesrestoration.gov/content/documents/system_wide_ecological_indicators/2016_system_wide_ecological_indicators.pdf

Cairns J, McCormick PV, Niederlehner BR. 1993. A proposed framework for developing indicators of ecosystem health. *Hydrobiologia* 236:1–44.

Carter GA, Lucas KL, Biber PD, Criss GA, Blossom GA. 2011. Historical changes in seagrass coverage on the Mississippi barrier islands, northern Gulf of Mexico, determined from vertical aerial imagery (1940–2007). *Geocarto Int* 26(8):663–673.

[CBBEP] Coastal Bend Bays & Estuaries Program. 2010. Environmental indicators report. April 2010. Corpus Christi (TX). 44 p. [accessed 2019 Mar 11]. <http://www.cbbep.org>

[CCME] Canadian Council of Ministers of the Environment. 1996. A framework for ecological risk assessment: General guidance. PN 1195. Ottawa (ON): Environment Canada. 32 p.

[CEC] Commission for Environmental Cooperation. 2011. A guide to ecological scorecards for marine protected areas in North America. Montreal (QC): North American Commission for Environmental Cooperation. 50 p. [accessed 2019 Mar 11]. www3.cec.org/islandora/en

[CERCLA] Comprehensive Environmental Response, Compensation and Liability Act of 1980. 1980. 43 C.F.R. 11.14. [accessed 2019 Mar 11]. <https://www.govinfo.gov>

Cormier SM, Smith M, Norton S, Nieheisel T. 2000. Assessing ecological risks in watersheds: a case study of problem formulation in the Big Darby Creek Watershed, Ohio, USA. *Environ Toxicol Chem* 19:1082–1096.

Couvillion BR, Barras JA, Steyer GD, Sleavin W, Fischer M, Beck H, Trahan N, Griffin B, Heckman D. 2011. Land area change in coastal Louisiana from 1932 to 2010. Pamphlet to accompany Scientific Investigations Map 3164. Reston (VA): US Geological Survey.

Dale VH, Beyeler SC. 2001. Challenges in the development and use of ecological indicators. *Ecol Indic* 1:3–10.

Doren RF, Trexler JC, Gottlieb AD, Harwell MC. 2009. Ecological indicators for system-wide assessment of the greater Everglades ecosystem restoration program. *Ecol Indic* 9S:S2–S16.

[EEA] European Environmental Agency. 1999. Environmental indicators: Typology and overview. EEA Technical Report no. 25. Copenhagen (DK). 19 p.

Egoh B, Drakou EG, Dunbar MB, Maes J, Willemsen L. 2012. Indicators for mapping ecosystem services: A review. European Commission. EUR 25456 – Joint Research Centre – Institute for Environment and Sustainability. Luxembourg (LU): Publications Office of the European Union. 111 p.

Environment Canada. 1994. A framework for ecological risk assessment at contaminated sites in Canada: Review and recommendations. Scientific Series No. 199. Ecosystem Conservation Directorate, Ottawa, Ontario, Canada. 108 p.

Felder DL, Camp DK, Tunnell JW Jr. 2009. An introduction to Gulf of Mexico biodiversity assessment. In: Felder DL, Camp DK, editors. Gulf of Mexico origin, waters and biota: Vol 1 Biodiversity. College Station (TX): Texas A&M Univ. p 1–12.

[GCERC] Gulf Coast Ecosystem Restoration Council. 2016. Comprehensive plan. Restoring the Gulf Coast's ecosystem and economy. New Orleans (LA). 31 p. [accessed 2019 Mar 11]. <https://www.restorethegulf.gov/>

Gentile JH, Harwell MA. 1998. The issue of significance in ecological risk assessments. *Hum Ecol Risk Assess* 4(4):815–828.

Gentile JH, Harwell MA, Cropper Jr WP Jr, Harwell CC, DeAngelis D, Davis S, Ogden JC, Lirman D. 2001. Ecological conceptual models: A framework and case study on ecosystem management for South Florida. *Sci Total Environ* 274:231–253.

Gentile JH, Harwell MA, van der Schalie W, Norton S, Rodier D. 1993. Ecological risk assessment: A scientific perspective. *J Hazard Mater* 35:241–253.

Gentile JH, Slimak MW. 1990. Endpoints and indicators in ecological risk assessment. In: MacKenzie DH, Hyatt DE, McDonald VJ, editors. Ecological indicators. Vol 2. New York (NY): Elsevier Applied Science. p 1385–1397.

- Goksøyr A, Förlin L. 1992. The cytochrome P-450 system in fish, aquatic toxicology and environmental monitoring. *Aquat Toxicol* 22(4):287–311.
- Gulf Coast Ecosystem Restoration Task Force. Executive Order 13554, 5 October 2010. 3 CFR 13554, p 258–262. Code of Federal Regulations, Washington (DC).
- Halpern BS, Longo C, Hardy D, McLeod KL, Samhouri JF, Katona SK, Kleisner K, Lester SE, O'Leary J, Ranelletti M et al. 2012. An index to assess the health and benefits of the global ocean. *Nature* 488:615–620 with Suppl.
- Harwell MA, Gentile JH. 2014. Assessing sea otter risks and the Exxon Valdez oil spill: New scenarios, attributable risk, and recovery. *Hum Ecol Risk Assess* 20(4):889–916. [accessed 2019 Mar 11]. <https://www.tandfonline.com/doi/abs/10.1080/10807039.2013.828513>
- Harwell MA, Gentile JH, Bartuska A, Harwell CC, Myers V, Obeysekera J, Ogden JC, Tosini SC. 1999. A science-based strategy for ecological restoration in South Florida. *Urban Ecosyst* 3:201–222.
- Harwell MA, Gentile JH, Cummins KW, Highsmith RC, Hilborn R, McRoy CP, Parrish J, Weingartner T. 2011. A conceptual model of natural and anthropogenic drivers and their influence on the Prince William Sound, Alaska, ecosystem. *Hum Ecol Risk Assess* 16:672–726. [accessed 2019 Mar 11]. <https://www.tandfonline.com/doi/full/10.1080/10807039.2010.501011>
- Harwell MA, Gentile JH, Parker KR. 2013. Characterizing ecological risks, significance, and recovery. In: Wiens JA, editor. Oil in the environment. Legacies and lessons of the Exxon Valdez oil spill. Cambridge (UK): Cambridge Univ. p 383–419.
- Harwell MA, Long JF, Bartuska AM, Gentile JH, Harwell CC, Myers V, Ogden JC. 1996. Ecosystem management to achieve ecological sustainability: The case of South Florida. *Environ Manage* 20:497–521.
- Harwell MA, Myers V, Young T, Bartuska A, Gassman N, Gentile JH, Harwell CC, Appelbaum S, Barko J, Causey B, Johnson C, McLean A, Smola R, Templet P, Tosini S. 1999. A framework for an ecosystem integrity report card. *BioScience* 49:543–556.
- Hattam C, Atkins JP, Beaumont N, Börger T, Böhne-Henrichs A, Burdon D, de Groot R, Hoefnagel E, Nunes PALD, Piwowarczyk J et al. 2015. Marine ecosystem services: Linking indicators to their classification. *Ecol Indic* 49:61–75.
- Holling CS. 1973. Resilience and stability of ecological systems. *Annu Rev Ecol Syst* 4:1–23.
- Horton KG, Van Doren BM, La Sorte FA, Cohen EB, Clipp HL, Buler JJ, Fink D, Kelly JF, Farnsworth A. 2019. Holding steady: Little change in intensity or timing of bird migration over the Gulf of Mexico. *Glob Change Biol* 25:1106–1118. [accessed 2019 Mar 11]. doi: 10.1111/gcb.14540
- Hunsaker CT, Carpenter DE, editors. 1990. Ecological indicators for the Environmental Monitoring and Assessment Program. Research Triangle Park (NC): USEPA Office of Research & Development. 429 p. EPA/600/3-90/060
- Hunsaker CT, Levine DA, Timmins SP, Jackson BL, O'Neill RV. 1990. Landscape characterization for assessing regional water quality. In: MacKenzie DH, Hyatt DE, McDonald VJ, editors. Ecological indicators. Vol 2. New York (NY): Elsevier. p 997–1006.
- [IAN] Integration & Application Network. 2007. 2006 Chesapeake Bay report card. Cambridge (MD): Center for Environmental Studies, Univ Maryland. [accessed 2019 Mar 11]. <http://ian.umces.edu/ecocheck/report-cards/chesapeake-bay/2006/>
- [IAN] Integration & Application Network. 2013. 2012 Chesapeake Bay report card. Cambridge (MD): Center for Environmental Studies, Univ Maryland. [accessed 2019 Mar 11]. <http://ian.umces.edu/ecocheck/report-cards/chesapeake-bay/2012/>
- Joye SB. 2015. Deepwater Horizon, 5 years on. *Science* 349(6248):592–593.
- Karr JR. 1981. Assessment of biotic integrity using fish communities. *Fisheries* 6:21–27.
- Kelble CR, Loomis DK, Lovelace S, Nuttle WK, Ortner PB, Fletcher P, Cook GS, Lorenz JJ, Boyer JN. 2013. The EBM-DPSER conceptual model: integrating ecosystem services into the DPSIR framework. *PLoS One* 8(8):e70766.
- Kelly JR, Harwell MA. 1989. Indicators of ecosystem response and recovery. Chapter 2. In: Levin SA, Harwell MA, Kelly JR, Kimball KD, editors. Ecotoxicology: Problems and approaches. New York (NY): Springer-Verlag. p 9–35.
- Kelly JR, Harwell MA. 1990. Indicators of ecosystem recovery. *Environ Manage* 15:527–545.
- Kujawinski EB, Kido Soule MC, Valentine DL, Boysen AK, Longnecker K, Redmond MC. 2011. Fate of dispersants associated with the Deepwater Horizon oil spill. *Environ Sci Technol* 45:1298–1306.
- Landres PB, Verner J, Thomas JW. 1988. Ecological uses of vertebrate indicator species: A critique. *Conserv Biol* 2:316–328.
- Mabus R. 2010. America's Gulf Coast. A long term recovery plan after the Deepwater Horizon oil spill. Report to the Gulf Coast Ecosystem Restoration Task Force. Gulf Coast Restoration Council, New Orleans, LA, USA. 127 p. [accessed 2019 Mar 11]. www.restorethegulf.gov
- MacKenzie DH, Hyatt DE, McDonald VJ, editors. 1990. Ecological indicators. Vol 1 and 2. New York (NY): Elsevier Applied Science. 1719 p.
- McKinney LD, Tunnell JW Jr, Harwell MA, Gentile JH, Dennison WC, Kelsey RH, Onuf C, Pollack JB, Stunz G, Withers K. 2018a. Texas Coast EcoHealth Metrics framework and prototype report card 2017. Corpus Christi (TX): Harte Research Institute for Gulf of Mexico Studies, Texas A&M University-Corpus Christi. [accessed 2019 Mar 11]. <https://www.hartheresearchinstitute.org/project/gulf-mexico-report-card-prototype-texas>
- McKinney LD, Tunnell JW Jr, Harwell MA, Gentile JH, Dennison WC, Kelsey RH, Onuf C, Pollack JB, Stunz G, Withers K. 2018b. Texas Coast EcoHealth Metrics framework technical support document 2017. 29 p. Corpus Christi (TX): Harte Research Institute for Gulf of Mexico Studies, Texas A&M University-Corpus Christi. [accessed 2019 Mar 11]. <https://www.hartheresearchinstitute.org/project/gulf-mexico-report-card-prototype-texas>
- Mills KE. 2006. A strategy for Gulf of Maine ecosystem indicators and state of the environment reporting. Gulf of Maine Ecosystem Indicator Partnership, Gulf of Maine Council on the Marine Environment, Buxton (ME). 56 p. [accessed 2019 Mar 11]. <http://www.gulfofmaine.org/2/esip-homepage>
- Mission-Aransas Reserve. 2019. Mission-Aransas National Estuarine Research Reserve. [accessed 2019 Mar 11]. <https://misionaransas.org>
- Morton RA, Bernier JC, Barras JA, Ferina NF. 2005. Rapid subsidence and historical wetland loss in the Mississippi Delta Plain: Likely causes and future implications. Open File Report 2005-1216. Reston (VA): US Department of the Interior, US Geological Survey. 124 p. [accessed 2019 Mar 11]. <http://pubs.usgs.gov/of/2005/1216/ofr-2005-1216.pdf>
- [NAS] National Academy of Sciences. 2012. Approaches for ecosystem services valuation for the Gulf of Mexico after the Deepwater Horizon oil spill. Interim report. Committee on the Effects of the Deepwater Horizon Mississippi Canyon–252 oil spill on ecosystem services in the Gulf of Mexico. Washington (DC): National Research Council, National Academies Pr. 162 p.
- NatureServe. 2019. NatureServe Vista. A powerful scenario-based assessment and planning tool. NatureServe, Arlington (VA). [accessed 2019 Mar 11]. <http://www.natureserve.org/conservation-tools/natureserve-vista>
- Newstead D, Fitzsimmons O. 2017. Post-Harvey Mid-Coast Rookery Island preliminary damage report. Report to the Coastal Bend Bay & Estuaries Program, 14 September 2017. Coastal Bend Bays & Estuaries Program, Corpus Christi, TX, USA. 15 p. [accessed 2019 Mar 11]. <https://www.gulfbase.org/project/indicators-and-assessment-framework-ecological-health-and-ecosystem-services>
- [NOAA] National Oceanic and Atmospheric Administration. 1996a. Injury assessment: Guidance document for Natural Resource Damage Assessment under the Oil Pollution Act of 1990. Damage Assessment and Restoration Program, National Oceanic and Atmospheric Administration, Silver Spring (MD). 214 p. [accessed 2019 Mar 11]. <https://darrp.noaa.gov/sites/default/files/Injury%20assessment.pdf>
- [NOAA] National Oceanic and Atmospheric Administration. 1996b. Restoration planning: Guidance document for Natural Resource Damage Assessment under the Oil Pollution Act of 1990. Damage Assessment and Restoration Program, National Oceanic and Atmospheric Administration, Silver Spring (MD). 169 p. [accessed 2019 Mar 11]. <https://darrp.noaa.gov/sites/default/files/Restoration%20Planning.pdf>
- [NOAA] National Oceanic and Atmospheric Administration. 2010. Oil Pollution Act of 1990 (OPA) OPA guidance. Damage Assessment, Remediation, and Restoration Program (DARRP), NOAA, Silver Spring (MD). 1 p. [accessed 2019 Mar 11]. <https://darrp.noaa.gov/oil-pollution-act-opa-1990>
- [NOAA] National Oceanic and Atmospheric Administration. 2011. Florida Keys National Marine Sanctuary condition report 2011. NOAA Silver

- Spring (MD): Office of National Marine Sanctuaries. 105 p. [accessed 2019 Mar 11]. http://sanctuaries.noaa.gov/science/condition/pdfs/fknms_highres.pdf
- [NOAA] National Oceanic and Atmospheric Administration. 2 Aug 2017. Gulf of Mexico 'dead zone' is the largest ever measured. *NOAA News and Features*. NOAA Silver Spring (MD) 1 p. [accessed 2019 Mar 11]. <https://www.noaa.gov/media-release/gulf-of-mexico-dead-zone-is-largest-ever-measured>
- Odum EP. 1969. The strategy of ecosystem development. *Science* 164: 262–270.
- Odum EP. 1985. Trends expected in stressed ecosystems. *BioScience* 35:419–422.
- [OECD] Organisation for Economic Co-operation and Development. 1991. Environmental indicators: A preliminary set. Paris (FR). 80 p.
- [OECD] Organisation for Economic Co-operation and Development. 1993. OECD core set of indicators for environmental performance reviews. Paris (FR). Environment Monographs No. 83. 39 p.
- Ogden JC, Davis SM, Barnes T, Jacobs KJ, Gentile JH. 2005. Total system conceptual ecosystem model. *Wetlands* 25:955–979.
- Ogden JC, Davis SM, Jacobs KJ, Barnes T, Fling HE. 2005. The use of conceptual ecological models to guide ecosystem restoration in South Florida. *Wetlands* 25:795–809.
- [OPA 90] Oil Pollution Act of 1990. 1990. 15 C.F.R. 990.30. [accessed 2019 Mar 11]. <https://www.govinfo.gov/>
- [OSTP] Office of Science and Technology Policy. 1999. Ecological risk assessment in the federal government. Washington (DC): Committee on Environment and Natural Resources of the National Science and Technology Council. CERN/5-99/001. 219 p.
- Pantus FJ, Dennison WC. 2005. Quantifying and evaluating ecosystem health: A case study from Moreton Bay, Australia. *Environ Manage* 36:757–771.
- Rabalais NN, Turner RE, Scavia D. 2002. Beyond science into policy: Gulf of Mexico hypoxia and the Mississippi River. *BioScience* 52:129–142.
- Rapport DJ, Regier HA, Hutchinson TC. 1985. Ecosystem behavior under stress. *Am Nat* 125:617–640.
- [RESTORE Act] Resources and Ecosystems Sustainability, Tourist Opportunities, and Revived Economies of the Gulf Coast States Act of 2012. 2012. Subtitle F – Gulf Coast Restoration. [accessed 2019 Mar 11]. <https://www.restorethegulf.gov/sites/default/files/RESTORE%20ACT%20July2012.pdf>. 20 p. and RESTORE Act Regulations for the Gulf Coast Restoration Trust Fund 31 C.F.R. 34 FR 80(239)77239–77252. 2015. [accessed 2019 Mar 11]. https://www.treasury.gov/services/restore-act/Documents/Final%20Rule_Federal%20Register_2015-31431.pdf
- Stanzel K. 2017. Rookery island suitability analysis. Green Wing Environmental, LLC, draft report submitted to Mission-Aransas Reserve, October 2017. 52 p. [accessed 2019 Mar 11]. <https://www.gulfbase.org/project/indicators-and-assessment-framework-ecological-health-and-ecosystem-services>
- Stephan CE, Mount DI, Hansen DJ, Gentile JH, Chapman GA, Brungs WA. 1985. Guidelines for deriving numerical national water quality criteria for the protection of aquatic organisms and their uses. Washington (DC): USEPA Office of Research & Development, Environmental Research Laboratories. USEPA PB85-227048. 59 p. [accessed 2019 Mar 11]. <http://water.epa.gov/scitech/swguidance/standards/criteria/aqlife/index.cfm#final>
- Stunz G, Hall Q, Francis J. 2018. Fisheries health support data. In: McKinney LD, Tunnell JW Jr, Harwell MA, Gentile JH, Dennison WC, Kelsey RH, Onuf C, Pollack JB, Stunz G, Withers K, editors. 2018. Texas Coast EcoHealth Metrics Framework technical support document 2017. Corpus Christi (TX): Harte Research Institute for Gulf of Mexico Studies, Corpus Christi (TX): Harte Research Institute for Gulf of Mexico Studies, Texas A&M University-Corpus Christi. 25 p. [accessed 2019 Mar 11]. <https://www.hartheresearchinstitute.org/project/gulf-mexico-report-card-prototype-texas>
- Suter GW II. 2007. Ecological risk assessment. Boca Raton (FL): CRC. 643 p.
- [TWCS] Texas Colonial Waterbird Society. 2019. Partnership: Texas Colonial Waterbird Society. A coalition working to monitor, promote research and inform management of colonial waterbird populations in Texas. [accessed 2019 Mar 11]. https://tpwd.texas.gov/huntwild/wild/wildlife_diversity/twcs/
- The Bay Institute. 2003. Ecological scorecard. San Francisco Bay Index 2003. Novato (CA): The Bay Institute of San Francisco. 78 p. [accessed 2019 Mar 11]. https://bayecotarium.org/wp-content/uploads/scorecard_report.pdf
- [UNEP WCMP] United Nations Environment Programme World Conservation Monitoring Centre. 2011. Developing ecosystem service indicators: Experiences and lessons learned from sub-global assessments and other initiatives. Montréal (QC): Secretariat of the Convention on Biological Diversity. Technical Series No. 58. 118 p.
- [USEPA] US Environmental Protection Agency. 1992. Framework for ecological risk assessment. Washington (DC): USEPA Risk Assessment Forum. EPA/630/R-92/001. 41 p.
- [USEPA] US Environmental Protection Agency. 1998. Guidelines for ecological risk assessment. Washington (DC): USEPA Risk Assessment Forum. EPA/630/R-95/002F. 188 p.
- [USEPA] US Environmental Protection Agency. 2011. Gulf of Mexico regional ecosystem restoration strategy. Gulf Coast Restoration Task Force. December 2011. Gulf Coast Ecosystem Restoration Council, New Orleans (LA). [accessed 2019 Mar 11]. http://archive.epa.gov/gulfcoasttaskforce/web/pdf/gulfcoastreport_full_12-04-508-1.pdf. 120 p.
- [USEPA] US Environmental Protection Agency. 2012. National coastal condition report IV. Washington (DC): Office of Research and Development/Office of Water. EPA-842-R-10-003. [accessed 2019 Mar 11]. http://water.epa.gov/type/oceb/assessmonitor/nccr/upload/0_NCCR_4_Report_508_bookmarks.pdf. 334 p.
- [USEPA SAB] US Environmental Protection Agency Science Advisory Board. 2002. A framework for assessing and reporting on ecological condition: An SAB report. Washington (DC): Science Advisory Board. EPA-SAB-EPEC-02-009. 133 p.
- Weber J-L. 2010. Merging the ecosystem approach with the conventional PSR/DPSIR framework (Draft for discussion). New York (NY): Department of Economic and Social Affairs, Statistics Division, United Nations. ESA/STATISTICS/AC.228, EGM-FDES/1/16. 7 p.
- Williams MR, Longstaff BJ, Buchanan C, Llanso R, Dennison WC. 2009. Development and evaluation of a spatially-explicit index of Chesapeake Bay health. *Mar Pollut Bull* 59:14–25.
- Williams MR, Finoso S, Longstaff BJ, Dennison WC. 2010. Long-term trends in water quality and biotic metrics in Chesapeake Bay: 1986–2008. *Estuaries Coasts* 33:1279–1299.
- Withers K. 2018. Texas EcoHealth Metrics bird populations data support document. In: McKinney LD, Tunnell JW Jr, Harwell MA, Gentile JH, Dennison WC, Kelsey RH, Onuf C, Pollack JB, Stunz G, Withers K, editors. 2017. Texas Coast EcoHealth Metrics Framework technical support document 2017. Corpus Christi (TX): Harte Research Institute for Gulf of Mexico Studies, Texas A&M University-Corpus Christi. 35 p. [accessed 2019 Mar 11]. <https://www.hartheresearchinstitute.org/project/gulf-mexico-report-card-prototype-texas>